

# Chapter 4-Carbon and the Molecular Diversity of Life

## Objectives:

1. Explain the difference between **vitalism** and **mechanism**.
2. Explain how carbon's electron configuration accounts for its ability to form large, complex, and diverse organic molecules.
3. Describe how carbon skeletons may vary, and explain how this variation contributes to the diversity and complexity of organic molecules.
4. Describe the basic structure of a hydrocarbon and explain why these molecules are hydrophobic.
5. Distinguish among the three types of isomers: structural, geometric, and enantiomer.

## Misconceptions:

1. Students often misunderstand the interaction of hydrophobic molecules and water. Many students think that individual oil and water molecules repel each other. Explain to students that individual hydrocarbon molecules are attracted to water molecules, but with a force much less than the attraction of water molecules to each other.
2. Students find it difficult to understand the differences among structural isomers, geometric isomers, and enantiomers. Three-dimensional models, or pictures of such models, are very useful tools in discussion of these terms.

## Notes:

### **A. Carbon—The Backbone of Biological Molecules**

- Although cells are 70–95% water, the rest consists mostly of carbon-based compounds.
- Carbon is unparalleled in its ability to form large, complex, and diverse molecules.
- Carbon accounts for the diversity of biological molecules and has made possible the great diversity of living things.
- Proteins, DNA, carbohydrates, and other molecules that distinguish living matter from inorganic material are all composed of carbon atoms bonded to each other and to atoms of other elements.
- These other elements commonly include hydrogen (H), oxygen (O), nitrogen (N), sulfur (S), and phosphorus (P).

### ***1. Organic chemistry is the study of carbon compounds.***

- The study of carbon compounds, **organic chemistry**, deals with any compound with carbon (organic compounds).
- Organic compounds can range from simple molecules, such as CO<sub>2</sub> or CH<sub>4</sub>, to complex molecules such as proteins, which may weigh more than 100,000 daltons.
- The overall percentages of the major elements of life (C, H, O, N, S, and P) are quite uniform from one organism to another.
- However, because of carbon's versatility, these few elements can be combined to build an inexhaustible variety of organic molecules.
- Variations in organic molecules can distinguish even between individuals of a single species.
- The science of organic chemistry began in attempts to purify and improve the yield of products obtained from other organisms.
- Initially, chemists learned to synthesize simple compounds in the laboratory, but had no

success with more complex compounds.

- The Swedish chemist Jons Jacob Berzelius was the first to make a distinction between organic compounds that seemed to arise only in living organisms and inorganic compounds that were found in the nonliving world.
- This led early organic chemists to propose **vitalism**, the belief that physical and chemical laws did not apply to living things.
- Support for vitalism began to wane as organic chemists learned to synthesize complex organic compounds in the laboratory.
- In the early 1800s, the German chemist Friedrich Wöhler and his students were able to synthesize urea from totally inorganic materials.
- In 1953, Stanley Miller at the University of Chicago set up a laboratory simulation of chemical conditions on the primitive Earth and demonstrated the spontaneous synthesis of organic compounds.
- Such spontaneous synthesis of organic compounds may have been an early stage in the origin of life.
- Organic chemists finally rejected vitalism and embraced **mechanism**, accepting that the same physical and chemical laws govern all natural phenomena including the processes of life.
- Organic chemistry was redefined as the study of carbon compounds regardless of their origin.
- Organisms do produce the majority of organic compounds.
- The laws of chemistry apply to inorganic and organic compounds alike.

## ***2. Carbon atoms can form diverse molecules by bonding to four other atoms.***

- With a total of 6 electrons, a carbon atom has 2 in the first electron shell and 4 in the second shell.
- Carbon has little tendency to form ionic bonds by losing or gaining 4 electrons to complete its valence shell.
- Instead, carbon usually completes its valence shell by sharing electrons with other atoms in four covalent bonds.
- This **tetravalence** by carbon makes large, complex molecules possible.
- When carbon forms covalent bonds with four other atoms, they are arranged at the corners of an imaginary tetrahedron with bond angles of  $109.5^\circ$ .
- In molecules with multiple carbons, every carbon bonded to four other atoms has a tetrahedral shape.
- However, when two carbon atoms are joined by a double bond, all bonds around those carbons are in the same plane and have a flat, three-dimensional structure.
- The three-dimensional shape of an organic molecule determines its function.
- The electron configuration of carbon makes it capable of forming covalent bonds with many different elements.
- The valences of carbon and its partners can be viewed as the building code that governs the architecture of organic molecules.
- In carbon dioxide, one carbon atom forms two double bonds with two different oxygen atoms.
- In the structural formula,  $\text{O}=\text{C}=\text{O}$ , each line represents a pair of shared electrons. This arrangement completes the valence shells of all atoms in the molecule.
- While  $\text{CO}_2$  can be classified as either organic or inorganic, its importance to the living world is clear.
- $\text{CO}_2$  is the source of carbon for all organic molecules found in organisms. It is usually fixed into organic molecules by the process of photosynthesis.

- Urea,  $\text{CO}(\text{NH}_2)_2$ , is another simple organic molecule in which each atom forms covalent bonds to complete its valence shell.

### ***3. Variation in carbon skeletons contributes to the diversity of organic molecules.***

- Carbon chains form the skeletons of most organic molecules.
- The skeletons vary in length and may be straight, branched, or arranged in closed rings.
- The carbon skeletons may include double bonds.
- Atoms of other elements can be bonded to the atoms of the carbon skeleton.
- **Hydrocarbons** are organic molecules that consist of only carbon and hydrogen atoms.
- Hydrocarbons are the major component of petroleum, a fossil fuel that consists of the partially decomposed remains of organisms that lived millions of years ago.
- Fats are biological molecules that have long hydrocarbon tails attached to a nonhydrocarbon component.
- Petroleum and fat are hydrophobic compounds that cannot dissolve in water because of their many nonpolar carbon-to-hydrogen bonds.
- **Isomers** are compounds that have the same molecular formula but different structures and, therefore, different chemical properties.
- For example, butane and isobutane have the same molecular formula,  $\text{C}_4\text{H}_{10}$ , but butane has a straight skeleton and isobutane has a branched skeleton.
- The two butanes are **structural isomers**, molecules that have the same molecular formula but differ in the covalent arrangement of atoms.
- **Geometric isomers** are compounds with the same covalent partnerships that differ in the spatial arrangement of atoms around a carbon–carbon double bond.

## **B. Functional Groups:**

### **Objectives:**

1. Name the major functional groups found in organic molecules. Describe the basic structure of each functional group and outline the chemical properties of the organic molecules in which they occur.

### **Notes:**

#### ***1. Functional groups are involved in chemical reactions and contribute to the molecular diversity of life.***

- The components of organic molecules that are most commonly involved in chemical reactions are known as **functional groups**.
- If we consider hydrocarbons to be the simplest organic molecules, we can view functional groups as attachments that replace one or more of the hydrogen atoms bonded to the carbon skeleton of the hydrocarbon.
- Each functional group behaves consistently from one organic molecule to another.
- The number and arrangement of functional groups help give each molecule its unique properties.
- As an example, the basic structure of testosterone (a male sex hormone) and estradiol (a female sex hormone) is the same.
- Both are steroids with four fused carbon rings, but they differ in the functional groups attached to the rings.
- These functional groups interact with different targets in the body.
- There are six functional groups that are most important to the chemistry of life: hydroxyl,

carbonyl, carboxyl, amino, sulfhydryl, and phosphate groups.

- All are hydrophilic and increase the solubility of organic compounds in water.
- In a **hydroxyl** group ( $\text{—OH}$ ), a hydrogen atom forms a polar covalent bond with an oxygen atom, which forms a polar covalent bond to the carbon skeleton.
- Because of these polar covalent bonds, hydroxyl groups increase the solubility of organic molecules.
- Organic compounds with hydroxyl groups are alcohols, and their names typically end in **-ol**.
- A **carbonyl** group ( $\text{>CO}$ ) consists of an oxygen atom joined to the carbon skeleton by a double bond.
- If the carbonyl group is on the end of the skeleton, the compound is an **aldehyde**.
- If the carbonyl group is within the carbon skeleton, then the compound is a **ketone**.
- Isomers with aldehydes versus ketones have different properties.
- A **carboxyl** group ( $\text{—COOH}$ ) consists of a carbon atom with a double bond to an oxygen atom and a single bond to the oxygen of a hydroxyl group.
- Compounds with carboxyl groups are **carboxylic acids**.
- A carboxyl group acts as an acid because the combined electronegativities of the two adjacent oxygen atoms increase the dissociation of hydrogen as an ion ( $\text{H}^+$ ).
- An **amino** group ( $\text{—NH}_2$ ) consists of a nitrogen atom bonded to two hydrogen atoms and the carbon skeleton.
- Organic compounds with amino groups are **amines**.
- The amino group acts as a base because the amino group can pick up a hydrogen ion ( $\text{H}^+$ ) from the solution.
- Amino acids, the building blocks of proteins, have amino and carboxyl groups.
- A **sulfhydryl** group ( $\text{—SH}$ ) consists of a sulfur atom bonded to a hydrogen atom and to the backbone.
- This group resembles a hydroxyl group in shape.
- Organic molecules with sulfhydryl groups are **thiols**.
- Two sulfhydryl groups can interact to help stabilize the structure of proteins.
- A **phosphate** group ( $\text{—OPO}_3^{2-}$ ) consists of a phosphorus atom bound to four oxygen atoms (three with single bonds and one with a double bond).
- A phosphate group connects to the carbon backbone via one of its oxygen atoms.
- Phosphate groups are anions with two negative charges, as two protons have dissociated from the oxygen atoms.
- One function of phosphate groups is to transfer energy between organic molecules.
- Adenosine triphosphate, or ATP, is the primary energy-transferring molecule in living cells.

## ***2. These are the chemical elements of life.***

- Living matter consists mainly of carbon, oxygen, hydrogen, and nitrogen, with smaller amounts of sulfur and phosphorus.
- These elements are linked by strong covalent bonds.
- Carbon, with its four covalent bonds, is the basic building block in molecular architecture.
- The great diversity of organic molecules with their special properties emerges from the unique arrangement of the carbon skeleton and the functional groups attached to the skeleton.

# **Key Terms**

adenosine triphosphate  
(ATP)  
amino group  
carbonyl group  
carboxyl group

enantiomer  
functional group  
geometric isomer  
hydrocarbon  
hydroxyl group

isomer  
organic chemistry  
phosphate group  
structural isomer  
sulfhydryl group

## Word Roots

**carb-** = coal (carboxyl group: a functional group present in organic acids, consisting of a carbon atom double-bonded to an oxygen atom and a hydroxyl group)

**enanti-** = opposite (enantiomer: molecules that are mirror images of each other)

**hydro-** = water (hydrocarbon: an organic molecule consisting only of carbon and hydrogen)

**iso-** = equal (isomer: one of several organic compounds with the same molecular formula but different structures and, therefore, different properties)

**sulf-** = sulfur (sulfhydryl group: a functional group that consists of a sulfur atom bonded to an atom of hydrogen)

**thio-** = sulfur (thiol: organic compounds containing sulfhydryl groups) .